

Chillerton and Gatcombe parish verge survey 2020

(This coding is optional and can be placed next to each plant identified)

F - Plant is frequently found. O - Plant is occasionally found. R - Plant is rarely found.

Name of surveyor(s) Carolyn and Steve Hoyle.	Date June 6th 2020
Place and length of verge Beginning at "Ramsdown" on Berry Shute, down the hill to the junction with Roslin Farm Lane (Approx' 400 - 450 metres)	
Verge type and orientation. South facing side of the lane is a shallow banked verge flanked by woodland and a small ditch and develops toward the Roslin end into a steeper bank flanked by farmland. The North facing bank is completely flanked by arable land but is separated by a deep ditch toward the lower end. The South facing verge has many species at the upper end, but where the verges are close to arable land (the lower South and North facing sides) they become much more overgrown with nettle, dock and bramble.	
Plants have finished flowering Not recorded.	
Plants in flower Dock (F) Crosswort (O) Greater Stitchwort (O) Red Campion (O) Ribwort Plantain (F) Germander Speedwell (O) Herb Robert (O) Sorrel (F) Hedge Bedstraw (R) Buttercup (O) Creeping Cinquefoil (O) Hogweed (F) Common Nettle (F) White Clover (O) Hemlock Dropwort (F) Brooklime (R) Thistle (O) Bramble (F) Dog Rose (pink) (R) Field Rose (White) (R) Woodsage (R) Cut-leaved Cranesbill (R) Meadow Vetchling (O) Hedge Woundwort(O) Smooth Hawk'sbeard (F) Sheeps Sorrel (R) Deadly Nightshade (R) Herb Bennet (O) Dwarf Elder (R)	
Plants with foliage or are about to flower Hazel (O) Bindweed (R) Fern/Bracken (R) Teasel (O)) Cleavers (O) Rosebay Willowherb (R) Silverweed (O) Yarrow (O) Ragwort (R) Fleabane (R)	